

AD-A046 965

TEXAS UNIV HEALTH SCIENCE CENTER AT DALLAS  
METABOLISM OF KININS AND ANGIOTENSINS IN THE KIDNEY.(U)  
NOV 77 P E WARD, E G ERDOS  
TR-9

F/G 6/1

N00014-75-C-0807

NL

UNCLASSIFIED

| OF |

AD  
A046 965



END  
DATE  
FILMED

12-77

DDC

AD A 0 46965

12

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0807

TECHNICAL REPORT NO. 9

Metabolism of Kinins and Angiotensins in the Kidney

Prepared by

Ervin G. Erdős


For Publication in  
Kininogenases

Departments of Pharmacology and Internal Medicine  
University of Texas Health Science Center at Dallas  
5323 Harry Hines Boulevard  
Dallas, Texas 75235

04 November 1977

Reproduction in whole or in part is permitted for  
any purpose of the United States Government

Distribution of this report is unlimited



AD No. \_\_\_\_\_  
DDC FILE COPY

1. GOVT ACCESSION NO. 2. RECIPIENT'S CATALOG NUMBER Technical Report No. 9 ✓	
3. TITLE (and Subtitle) <b>METABOLISM OF KININS AND ANGIOTENSINS IN THE KIDNEY.</b>	
4. TYPE OF REPORT & PERIOD COVERED Technical 6/1/77 - 12/15/78	
5. PERFORMING ORG. REPORT NUMBER Patrick E. Ward and Ervin G. Erdős	
6. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0807	
7. PERFORMING ORGANIZATION NAME AND ADDRESS Univ. Tx. Health Sci. Center at Dallas Dept. of Pharmacology 5323 Harry Hines Blvd., Dallas, TX 75235	
8. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Biological & Medical Sciences Division Medical and Dental Sciences Program, Code 444 Arlington, Virginia	
9. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 9 Technical rept. 1 Jun 77-15 Dec 78	
10. SECURITY CLASS. (of this report) Unclassified	
11. DISTRIBUTION STATEMENT (of this Report) Distribution of this document is unlimited 12 12p. 14 TR-9	
12. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) 11 4 Nov 77	
13. SUPPLEMENTARY NOTES To be published in <u>Kininogenases</u> , G. L. Haberland (ed.) Schattauer Publ. Stuttgart.	
14. KEY WORDS (Continue on reverse side if necessary and identify by block number) Angiotensin, bradykinin, renal tubules, kallikrein, prostaglandins	
15. ABSTRACT (Continue on reverse side if necessary and identify by block number) The paper suggest a mechanism for the inactivation of kinins and angiotensins in the brush border of the nephron.	




**METABOLISM OF KININS AND ANGIOTENSINS IN THE KIDNEY**

**Patrick E. Ward and Ervin G. Erdős**

**Departments of Pharmacology and Internal Medicine  
University of Texas Health Science Center at Dallas  
5323 Harry Hines Blvd.  
Dallas, Texas 75235**

**Send Galley Proofs to:**

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	
	



The kidney is rich in the enzymes which form or inactivate kinins and angiotensins. These vasoactive peptides have potent effects on renal blood flow, salt-water balance and prostaglandin release. If intrarenal generations of these peptides are physiologically significant modifiers of renal function, their rates of inactivation are equally important.

The kidney has a very high kininase content (1). In the homogenized kidney most of this activity sediments with the microsomal fraction. Erdős and Yang (2) described several of these renal kininases. One of them is a carboxypeptidase (kininase I) while the other (kininase II) is identical with the angiotensin I converting enzyme (3,4).

Kinins (5) and angiotensins as well (6), are rapidly and nearly completely inactivated when infused into the kidney. Little or no intact bradykinin reaches the venous effluent or urine. Urine does, however, contain kinins (7), which are apparently released intrarenally.

Unlike renal kallikrein which is localized primarily in the cortex (8), renal kininase and converting enzyme are present in both cortex and outer medulla (9). After fractionation of renal microsomes into plasma membrane and endoplasmic reticulum-enriched fractions, renal kininase, converting enzyme (10) and angiotensinase, (possibly A type), were found predominantly in an enriched fraction of plasma membrane (11,12). Isolated brush borders of proximal tubules (Fig. 1) are particularly rich in these enzymes while little activity was found in isolated glomeruli (12,13). Transplanted renal tumors, thought to be derived from cells of the pars recta of proximal tubules, also contain kininase, converting enzyme and angiotensinase (12,14). In contrast, kallikrein was not present in the brush border of proximal tubules because it originates from distal tubules.

Another approach to show kininase II in the kidney consisted of using fluorescent antibody to purified swine kidney enzyme. By this technique converting



enzyme was found throughout the nephron, but more concentrated in the proximal tubule (14) than in the distal tubule (Fig. 2). In rabbits (15) converting enzyme is also present in the proximal tubules. Both studies reported much lower levels associated with the glomerular tufts (14, 15).

Using micropuncture of proximal and distal tubules, Carone et al. (16) and Pullman et al. (17) have shown that the proximal tubule is highly active in hydrolyzing added bradykinin and angiotensin II. Little or no activity was found in the distal tubule.

After considering all of these observations, we suggest the following mechanism for renal inactivation of kinins and angiotensins (Fig. 3). If kinins and angiotensins released intrarenally have important effects on water and sodium metabolism and prostaglandin release, the kidney must be capable of inactivating these peptides entering the nephron from extra-renal sources. The low molecular weight of these peptides makes it likely that any plasma kinin or angiotensin that escapes inactivation in the circulation is filtered at the glomerulus. High levels of kininase and angiotensinase on the huge surface of the brush border of the proximal tubules then ensure complete removal of these peptides from the tubular filtrate which eventually reaches the distal tubules. At the distal tubule, renal kallikrein (11,12,18) interacts with a plasma or tissue kininogen to release renal kinin. The level of kinin at this site can be closely controlled since all traces of extra-renal kinin have been removed. Kallikrein by liberation of (19), renal kinin (20) may alter ion transport, water reabsorption and liberate prostaglandins (21) in collecting tubules and ducts.



1. E. Werle, W. Götze, A. Kappler, *Biochem. Z.* 289, 217 (1937).
2. E. G. Erdős, H. Y. T. Yang, in *Hypotensive Peptides* (Eds. E. G. Erdős, N. Back and P. Sicuteri), p. 235.. Springer-Verlag, New York (1966).
3. E. G. Erdős and H. Y. T. Yang, *Life Sci.* 6, 569 (1967).
4. H. Y. T. Yang, E. G. Erdős and Y. Levin, *Biochim. Biophys. Acta* 214, 374 (1970).
5. K. Abe, *Tohoku. J. Exp. Med.* 87, 175 (1965).
6. J. G. Ledingham and W. P. Leary, in *Angiotensin* (Eds. I. H. Page and F. M. Bumpus), P. 111. Springer-Verlag, New York (1974).
7. I. Miwa, E. G. Erdős, and T. Seki, *Proc. Soc. Exp. Biol. Med.* 131, 768 (1969).
8. K. Nustad, *Br. J. Pharmacol.* 39, 87 (1970).
9. P. E. Ward and I. H. Mills, *J. Endocrinol.* 67, 60P (1975).
10. E. G. Erdős, *Am. J. Med.* 60, 749 (1976).
11. P. E. Ward, C. D. Gedney, R. M. Dowben and E. G. Erdős, *Biochem. J.* 151, 755 (1975).
12. P. E. Ward, E. G. Erdős, C. D. Gedney, R. M. Dowben and R. C. Reynolds, *Biochem. J.* 157, 643, (1976).



13. P. E. Ward, W. Schultz, R. C. Reynolds and E. G. Erdős, Lab. Invest. in press, (1977).
14. E. R. Hall, J. Kato, E. G. Erdős and G. Oshima, Life Sci. 18, 1299 (1976).
15. P. R. B. Caldwell, B. C. Seegal, K. C. Hsu, M. Das and R. L. Soffer, Science 191, 1050 (1976).
16. F. A. Carone, T. N. Pullman, S. Oparil and S. Nakamura, Am. J. Physiol. 230, 1420 (1976).
17. T. N. Pullman, S. Oparil and F. A. Carone, Am. J. Physiol. 226, 747 (1975).
18. T. B. Ørstavik, K. Nustad, P. Brandtzaeg and J. U. Pierce, J. Histochem. Cytochem. 24, 1037 (1976).
19. A. G. Scicli, O. A. Carretero, A. Hampton and N. B. Oza, Fed. Proc. Fed. Am. Soc. Exp. Biol. 34, 376 (1975).
20. A. G. Scicli, R. Gandolfi, O. A. Carretero, Fed. Proc. Fed. Am. Soc. Exp. Biol. 36, 1015 (1977).
21. J. C. McGiff, H. D. Itskovitz, A. Terragno and P. U. K. Wong, Fed. Proc. Fed. Am. Soc. Exp. Biol. 35, 175 (1976).



## Legends

Fig. 1. Scanning electron micrograph of brush border of proximal tubules in rat kidney (12).


Fig. 2 Fluorescent photomicrography (X90) of a 4  $\mu$ m section of swine kidney cortex incubated with fluorescein isothiocyanate labeled antibody against angiotensin I converting enzyme (14).



**Footnotes:**

**These studies have been supported in part by N.I.H., U.S.P.H.S. HL 14187,  
16320, 1F 22 HL 00643 and ONR N00014 75 0807.**







# RENAL METABOLISM OF KININS & ANGIOTENSINS




Figure 3